

## **Quality of life of psychiatric patients**

**Demerdash, A. El-Shafei, T. Esmaeil, R. Ragheb, K. Atteya, H. Hatata, H.**

**Current Psychiatry. Ain Shams University (Print)**

**2008 v15 n1**

Success is measured by social and mental well-being as well as by standard of living and quality of life, of the family as a whole and of each member. The consequences of psychiatric disorders have become a major focus of epidemiologic research over the past decade. Psychiatric disorders have substantial financial costs and broad social consequences for patients, their families and the larger society. However, important questions about the effects underlying these processes remain. The concept of quality of life has gained currency over the last twenty years as a way of assessing the global outcome of a variety of medical condition including psychiatric disorder. It has the advantage of generating an overview based on a number of more specific domains of functioning, activity and accesses to resources that together influence people's sense of well-being. The aim of this study was to explore the patient's quality of life. It was found that patients reported higher satisfaction toward work and activity, social relations, finance, work and school, low and safety and health.

While, the highest unsatisfaction with living situation and family. The current data provide evidence that, a statistical significant and positive correlation was found between patients with personality disorders and global rating, living situation, family relations, social relations, and work and activity. While there is statistical significant and negative correlation with low and safety and health (Significance at  $P < 0.05$ ). Patients with personality disorders showed marked unsatisfaction with their different category of quality of life. The results showed that family relations among quality of life of patients with different psychiatric disorders have statistical significant and negative correlations with depressive disorders, schizophrenia spectrum disorders and somatoform disorders ; while there is a statistical significant and positive correlations with bipolar disorders and anxiety disorders.