

Psychobiological Aspects of Patients with Lichen Planus

EL.Tawil M¹, Sediki N², Hassan H³

¹Department of Psychiatry, Faculty of Medicine, Cairo University, Cairo, Egypt

²Department of Dermatology, Faculty of Medicine, Cairo University, Cairo, Egypt

³Department of Chemical & Clinical Pathology, Faculty of Medicine, Cairo University, Cairo, Egypt

ABSTRACT

Background: Lichen planus, a psychocutaneous disorder, is associated with psychoneuroendocrine and psychoimmunological comorbidities. **Objective:** to assess psychiatric comorbidities in a group of patients with lichen planus and its impact on health related quality of life. Also we tried to assess some psychoneuroimmunological and pschoneuroendocrinological pathogenic mechanisms in lichen planus as a model in psychodermatoses. **Subjects and methods:** We included 70 patients with lichen planus All participants were subjected Psychological assessment using SF-36 health related quality of life, alexithymia (SAT9), Taylor manifest anxiety and Beck depression inventory (BDI), and laboratory measures including salivary cortisol and T cell subpopulations (CD+4 and CD8+). **Results:** Lichen plans patients showed higher frequency of psychiatric comorbidities (62.9%), poor health related quality of life and more frequent alexithymia (51.4%) Lichen planus patients with psychiatric comorbidities showed higher scoring on BDI, Taylor anxiety, poorer quality of life and higher tendency for endocrinological and immunological dysregulations. There were negative correlations between some variables and psychiatric comorbidity, quality of life, salivary cortisol and immunological dysregulations in patients with lichen planus. **Conclusion:** These findings may help dermatologists recognizing emotional factors in skin disorders. Consultation liaison psychiatrist may be useful in treating, ameliorating, improving quality of life and coping which may influence susceptibility to infections and immune response in a wide variety of psychodermatoses.

Key words: Lichen Planus, Psychobiological aspects, Egypt.

(Current Psychiatry 2009;16(4):370-80)

INTRODUCTION

Fyodor Desteovsky (1868) wrote that if you happen to have a wart on your nose or forehead, you cannot help imagining that no one in the world has anything else to do but stare at your wart, laugh at it and condemn you for it, even though you have discovered America.

Being a barrier between the external environment and the internal milieu, the skin plays several important functions both

physiological and psychological¹. The physiological role of the skin helps in the preservation of body homeostasis through protective barrier against external noxious stimuli and regulating body temperature. Psychologically, skin is an erogenous zone and channel for emotional discharge so that troubled skin could be a manifestation of unexpressed anger or an inner conflict due to external stress².

Brig et al. suggested that there is a diversionary symbiosis between skin and psyche³. Psychological stress could have a negative impact on healthy skin, exacerbating or precipitating dermatological disorders suggesting the presence of interface between psychiatry and dermatology⁴. In this way, dermatological disorders may be a skin language to express repressed feelings as symbols for crying out searching for love, respect and protection. Indeed high prevalence of psychiatric problems in patients with skin disorders was reported (20%-40%)⁵.

The interaction between mind and skin has led to the emergence of psychodermatology or psychocutaneous disorders concept. Psychodermatological or psychocutaneous disorders are classified into 4 categories⁶;

1. Psychophysiological (skin disease precipitated or exacerbated by stressful events).
2. Psychiatric disorders with dermatological symptoms (mostly occurring in somatoform disorders, anxiety, factitious, eating disorders)
3. Dermatologic disorders with psychiatric symptoms (emotional problems secondary to skin disease).
4. Miscellaneous; disorders or symptoms not otherwise classified (e.g.: cutaneous sensory pain, Gardener Diamond syndrome).

Dermatologic manifestation could be adverse effects from psychotropics (antipsychotics, antidepressants, anxiolytics and lithium) as well as changes in mood or behavior could result from steroids used to treat dermatological disorders⁷.

Lichen planus is one of the psychodermatological disorders affecting 1-2% of the general population⁸, occurring at any age (mean age of onset middle age), affecting both sexes with females being at higher risk for oral mucosal and genital involvement⁹. It is an obstinate disorder baffling not only the patients but also the practitioner. People with viral hepatitis or cirrhosis are at a high risk¹⁰.

Lichen planus is not an infectious disease. The cause is unknown, but it is classified as an autoimmune disorder which may be precipitated or exacerbated by psychosocial stressors¹¹. Patients with lichen planus experience stressful events before the onset of the disease⁹ Also higher level of anxiety and high salivary cortisol levels in patients with oral lichen planus¹².

Lichen planus manifested as itchy, reddish purple bumps affecting any part of the skin with predilection for inner sides of wrists, forearms, lower legs, genitalia, and to a lesser extent scalp, lower back or nails. Hypertrophic lichen planus is a variant of the disease appearing as thick, reddish-brown lesions covered with scales, mainly appearing on the shin⁸.

Lichen planus may affect oral mucous membranes with the risk of progression to oral squamous carcinoma¹³. Several studies reported an association between psychopathological disorder and lichen planus. Major life events especially illness or death of a dear one could precede or exacerbate coetaneous lichen planus¹¹.

The pathogenesis of lichen planus includes cell mediated immune mechanisms and neuroendocrine dysregulations mainly involving glucocorticoids¹⁴. Although there conflicting results regarding immunopathogenesis of lichen planus, the

major concept is that lichen planus is a T-cell mediated autoimmune disorder with a lymphotoxic processes against epidermal basal layer as an apoptotic mechanism¹⁵. Significant increase in CD8+ Tcell subpopulation was also noted¹⁶. The neuroendocrine mechanism may be mediated either centrally (HPA axis) or by independent cutaneous neuroendocrine axis¹⁷. Psychosocial stresses could exacerbate or precipitate lichen planus through neuroendocrine and neuroimmunologic mechanisms.

In a trial to investigate the interface between psychiatry and dermatology, our work aimed at:

- a. Screening for psychiatric manifestations in lichen planus.
- b. Comparing quality of life in lichen planus patients with and without psychiatric manifestations.
- c. Correlating some neuroendocrine and immune changes in lichen planus with psychiatric manifestations.

SUBJECTS AND METHODS

The study was conducted on 70 patients with coetaneous idiopathic (classic) lichen planus without oral mucosal involvement and 50 subjects with skin disorder with known negligible psychosomatic components as a control group, recruited from Asaad El-Hamid Center for Dermatology.

Inclusion criteria:

All participants in the study should fulfill the following criteria;

- 1- Age ranges 25-45 years.
- 2- Both sexes were included
- 3- At least 8 years of education

Exclusion criteria:

1. Age less than 25 or more than 45 years.
2. Organic mental syndrome or disorders.
3. Severe mental illnesses (schizophrenia, schizoaffective ...).
4. History of substance abuse disorders.
5. Chronic medical disorders.
6. Surgical operation in the previous 6 months prior to study
7. Medications with effects on immune system, behavior or mood at least 3 months prior to the study (Steroids, anti-inflammatory, antihistaminics, beta blockers, antibioticsetc.).
8. Pregnant women or women using oral or injectable contraceptives.

Methods

All participants in the study were subjected to the following after giving consent:

- 1- Complete medical history and thorough physical examination with special details regarding onset, duration, site and morphology of lichen planus lesions.
- 2- Psychiatric sheet.
- 3- ICD-10 Symptoms Checklist for mental disorders; a semi structured instrument intended for clinician assessment of the psychiatric symptoms and syndromes in F0-F6 categories of the International Classification of Diseases, Tenth Revision (18)
- 4- Psychometric tools;

a- Health related quality of life assessed by SF-36 instrument (HRQ). It consists of 8 subscales evaluating different domains (physical functioning, physical role, bodily pain, general health, social functioning, emotional role and mental health). Each domain scored from 0-100 with higher scores indicates better quality of life¹⁹.

b- Scored Archetypal Test of 9 elements (SAT9); to measure symbolic function in alexithymic individuals²⁰.

c- Beck Depression Inventory (BDI) a 21-item scale measuring symptoms related to cognitive, behavioral, affective and somatic components of depression²¹.

d- Taylor Manifests Anxiety Scale (TAMS) self reported scale consisting of 50 item statements indicative of anxiety from the Minnesota Multiphasic Personality Inventory (MMPI). Additional items have been added to control for social distractibility, lying and rigidity response²².

e- Gurmeet singhs presumptive life events stress scale. It contains 47 specific events common to individuals in a variety of situations plus the opportunity for the subject to list 3 other specific events that have occurred over the past year and then rates the positive or negative impact of these events.

Laboratory measures:

a- Salivary cortisol, measuring salivary cortisol has several advantages over blood cortisol including, stress free sampling, laboratory independence and low cost²³. Salivary cortisol is a measure of unbound cortisol fraction collected by salivette (sarstedt technique) and assayed by a technique combining automated HPLC (high performance liquid chromatography) and polarization fluorescence is used (24).

b- T lymphocytes subpopulations (CD4+ percentage and CD4+/CD8+ ratio) in peripheral blood. CD4+ (T-helper) and CD8+ (T-suppressor) are among T-lymphocytes subpopulations which state of the host immunity. Peripheral blood collected by venupuncture and flow

cytometer analysis (EPICS-XL) using dual fluorescence analysis.

Statistical analysis

Data were collected, computerized and revised for accuracy and consistency using soft ware package SPSS (version 11). Qualitative parameters assessed by frequency distribution, while means and standard deviation were used to describe quantitative parameters. Association between two qualitative variables was assessed using chi-square tests, while relation between quantitative variables was assessed using correlation analysis. Multiple statistical analyses were used to detect significant associations between different variables. Level of significance in all statistical tests was considered at $p < 0.05$.

RESULTS

This study included 70 patients with lichen planus and 50 control subjects. Both groups matched for age (34.32 ± 9.26 years in patients versus 32.97 ± 8.76 years in control group), sex (patient group; 51.5% females, 48.5% males, 52% females and 48% males in control group), marital status, educational, occupational and socioeconomic level.

Psychiatric symptoms as shown in table (1) showed statistically significant differences ($p < 0.005$) between patients with lichen planus and comparison group. Among lichen planus group phobic anxiety related symptoms were the most frequent (15.7% mixed anxiety depressives, 12.9% social phobia, followed by panic 11.45, obsessive compulsive 10%) while dysthymia was the least frequent (5.7%).

In table (2) mean scores of health related quality of life using short form 36 scale are compared in lichen planus and comparison group. Lower scoring indicating more negative influence on the quality of life were found in lichen planus patients but statistically significant differences ($p < 0.05$) could be revealed on five subscales of SF-36 namely physical functions, vitality, social function, emotional role limitation and mental health.

In table (3) lichen planus patients with psychiatric symptoms showed statistically significant lower mean scoring ($p < 0.05$) on bodily pain, vitality, social function, emotional role limitation and mental health subscales of SF-36 health related quality of life compared to lichen planus patients without psychiatric symptoms.

In table (4) Lichen planus group showed higher frequency to be alexithymic (51.4%) compared to (10%) of comparison group

Alexithymia showed statistically significant differences in patients with psychiatric symptoms compared to those without psychiatric symptoms as shown in table (5) 63.6% versus 30.8% respectively.

In table (6) higher scoring on Beck depression and Taylor anxiety scales were shown in lichen planus patients with psychiatric symptoms compared to lichen planus without psychiatric symptoms.

Table (7) demonstrated that lichen planus patients with psychiatric symptoms experienced more frequent undesirable events compared to those without psychiatric symptoms.

Table (1): ICD-10 symptoms checklist frequency in lichen planus & comparison groups.

ICD-10 symptom checklist	Lichen planus (N=70)		Comparison group (N=50)		p-value
	No	%	No	%	
Phobic anxiety					
Mixed anxiety and depressives	11	15.7	2	4	0.0001*
Social phobia	9	12.9	0	0	0.0001*
Panic	8	11.4	0	0	0.0001*
Obsessive compulsive (predominantly obsessional thoughts and ruminations)	7	10	1	2	0.0001*
Generalized anxiety	5	7.1	1	2	0.0002*
Mood (affective) disorders; dysthymia	4	5.7	0	0	0.0001*
No psychiatric disorder or manifestations	26	37.1	46	92	0.00001*

Table (2): Health related quality of life using SF-36 scale.

SF -36 Domain	Lichen planus (N=70)		Comparison group (N=50)		p-value
	Mean	# SD	Mean	#SD	
Physical function	50.82	32.51	75.19	21.19	0.001*
Physical role limitation	80.38	15.31	82.17	18.12	ns
Bodily pain	75.31	22.46	76.15	21.17	ns
General health	71.37	21.37	72.22	18.16	ns
Vitality	50.82	23.15	70.19	23.12	0.001*
Social function	50.14	25.12	80.19	24.13	0.0001*
Emotional role limitation	56.18	20.14	77.16	22.16	0.0001*
Mental health	57.16	22.13	76.19	31.12	0.0001*

* Significant p ($p < 0.05$)

Table (3): SF-36 quality of life scoring in lichen planus patients with and without psychiatric symptoms.

SF-Domain	Lichen planus with psychiatric symptoms (N=44)		Lichen planus without psychiatric symptoms (N=26)		p-value
	Mean	SD	Mean	SD	
Bodily pain	52.13	21.12	65.14	25.18	0.001*
Vitality	46.16	18.13	60.15	28.12	0.001*
Social function	48.38	26.19	58.14	21.13	0.0001*
Emotional role limitation	50.12	21.11	65.12	24.13	0.0001*
Mental health	44.16	21.13	60.12	23.19	0.001*

* Significant p (p<0.05).

Table (4): Frequency distribution of alexithymia using (SAT9) in lichen planus and comparison groups.

SAT9	Lichen planus (N=70)		Control group (N=50)		P-value
	No.	%	No.	%	
Alexithymics	36	51.4	5	10	0.0001*
Non-alexithymics	34	48.6	45	90	

*significant p (p<0.05)

Table (5): Frequency distribution of alexithymia in lichen planus with and without psychiatric symptoms.

SAT9	Lichen planus with psychiatric symptoms		Lichen planus without psychiatric symptoms		p-value
	No.	%	No.	%	
Alexithymics	28	63.6	8	30.8	0.0001*

* Significant p (p, 0.05)

Table (6): Mean scoring on beck depression inventory and Taylor manifest anxiety scales in lichen planus patients with and without psychiatric symptoms.

variable	Lichen planus with psychiatric symptoms		Lichen planus without psychiatric symptoms		P-value
	Mean	Sd	Mean	Sd	
Beck inventory	13.33	8.21	6.35	2.79	0.001*
Taylor anxiety	24.18	9.76	11.86	7.98	0.001*

* Significant p (p<0.05).

Table (7): Frequency of categorized events among lichen planus patients with and without psychiatric symptoms.

Categorized events	Lichen planus with psychiatric symptoms (N=44)		Lichen planus without psychiatric symptoms (N=26)		p-value
	No.	%	No.	%	
Undesirable	30	68.2	10	38.5	0.0001*
Desirable	14	31.8	16	61.5	0.001*

*Significant p (p<0.05).

Table (8): Frequency of salivary cortisol, CD4+ percent and CD4+/CD8+ ratio in lichen planus patients with and without psychiatric symptoms and control group.

Variable	Lichen with psychiatric symptoms (N=44)		Lichen without psychiatric symptoms (N=26)		Control group (N=50)		p
	N.	%	No.	%	No.	%	
Salivary cortisol (high level)	30	68.2	10	38.5	0	0	
Low CD4+ %	28	63.3	9	34.7	0	0	0.000*
Decreased CD4+/CD8+ ratio	30	68.2	11	42.3	0	0	

Table (9): Correlations between SF-quality of life, Taylor Anxiety scoring, salivary cortisol and immune dysregulations in lichen planus patients.

variable	SF-36 quality of life scoring	Taylor anxiety	High salivary cortisol	Immune changes
Morphology of Lesion (Hypertrophic)	-0.42*	-0.51*	-0.28*	-0.49*
Site of lesion; Genital	-0.25*	-0.38*	-0.39*	-0.41*
Upper limbs	-0.71*	-0.42*	-0.36*	-0.29*
Gender (females)	-0.35*	-0.33*	-0.29*	-0.37*
Occupation (professionals)	-0.68*	-0.63*		
Socioeconomic (low status)	-0.57*	-0.42*	-0.30*	-0.42*
Psychiatric symptoms	-0.69*	-0.71*	-0.48*	-0.52*

* Significant p (p<0.05).

Table (8) showed that lichen planus patients had increased salivary cortisol and immune dysregulations (low CD4+ % and low CD4+/CD8+ ratio) compared to control. Also, lichen planus patients with psychiatric symptoms showed statistically significant differences regarding neuroendocrine and immunological dysregulations compared to lichen planus patients without psychiatric symptoms (p<0.05).

Table (9) showed correlations between some variables (hypertrophic lesions, lesions affecting genitalia or upper limbs,

occupations as teachers', nurses' low economic status and presence of psychiatric symptoms) and poor quality of life, higher level of anxiety, high salivary cortisol and immune dysregulations

DISCUSSION

Psychodermatology is an interesting area for interface between psychiatry and dermatology, as there is a bidirectional interaction or crosstalk between the brain, psyche and skin. This crosstalk may stem

from the fact that the ectoderm is the embryologic origin for nervous and most skin tissue. In addition psychosocial stress could induce autoimmune or inflammatory skin disorders through neuroendocrine and neuroimmune dysregulations as well as dermatologic disorder may result in psychiatric morbidity and negative impact on quality of life. Having bad skin can result in poor self image, lack of confidence especially if the affected areas are exposed.

The present study showed that (63.6%) of lichen planus patients had psychiatrists symptoms with mixed anxiety and depressives being the most frequent (15.7%), followed by social phobia (12.9%), panic symptoms (11.4%), obsessive compulsive predominantly obsessional thoughts and ruminations (10%), and dystymia (5.7%) arranged in order. Magin et al. estimated that 30%-60% (nearly 10 fold that of the general population) of dermatological patients have psychiatric problems and dermatologists should not be surprised by that figure. This is not consistent with Mansur et al. who found that lichen planus patients did not differ from controls regarding psychiatric disorders⁹. This disagreement may be due to differences in methodology and sample selections.

Regarding health related quality of life our results showed statistically significant differences between lichen planus and control group on five subscales (physical function, vitality, social functions, emotional role and mental health). The presence of psychiatry symptoms in subgroup of lichen planus increases the negative impact on life quality as demonstrated by statistically significant lower scoring on SF-36 health related quality of life subscales in lichen planus

patients with psychiatric symptoms compared to those without psychiatric symptoms. Due to visibility of skin, dermatological disorder not only associated with cosmetic disfigurement but can result in a variety of psychopathological problems affecting patient, family and society²⁵. Some studies suggested that social success correlated positively with physical appearance and the more better looking you are, the more positively people will relate to you²⁶.

Our results showed alexithymia was more frequent in lichen planus than control group (51.4% versus 10%), and those with psychiatric symptoms had higher frequency of alexithymia than those without psychiatric symptoms with lichen planus (63.06% versus 30.8%) indicating that inhibited symbolic functions. This finding is consistent with Mansur et al. who reported statistically significant differences between lichen planus and controls⁹.

The results of this study showed that lichen planus patients with psychiatric symptoms scored more on anxiety symptoms using Taylor scale and BID than those without psychiatric symptoms. This may be due to social embarrassment, feeling of inadequacy, low self esteem and impaired coping strategies to deal with physical stress resulting from the disease and psychological stress. This finding is also in agreement with Mansur et al. who reported higher degree of anxiety in lichen planus compared to control⁹. Our results showed that lichen planus with psychiatric symptoms experienced more frequent undesirable events compared to those without psychiatric symptoms. This is consistent with many suggestions considering that psychosocial or stressful

events may precipitate or exacerbate dermatological disorders.

The results of this study showed statistically significant differences regarding neuroendocrine (salivary cortisol) and T-cell mediated immune dysregulations (decreased CD4+ % and low CD4+/CD8+ ratio) in lichen planus and control group and the presence of psychiatric symptoms is associated with more dysregulations. There are conflicting results regarding cortisol and cell mediated immune changes in lichen planus. The neuroendocrine and neuroimmunological findings in skin disorders may be difficult to interpret because it is not fully understood. There may be a multidirectional communications between skins, endocrine, immune, brain as well as other internal organs suggesting the existence of new layer for the regulation of global homeostasis²⁷. Some investigators hypothesized that skin has cutaneous neuroendocrine axes equivalent of the hypothalamic–pituitary–adrenal axis (HPA) that would regulate local response to stress independently from the central level¹⁷.

As an attempt to find patients with lichen planus are at higher risk to have poor quality of life, higher anxiety, endocrine and immune dysregulations; correlation analysis was done and we tried to explain these correlative relations between these variables and each other according to our study;

a- Lichen planus lesion morphology hypertrophic type. This may be explained by knowing that hypertrophic lesions tend to be more itchy sometimes severe enough to interfere with daily activities, hyperpigmentation, chronicity and scaly.

b- Lesion site ;1- genital lesion (may cause psychosexual discomfort, may be confused

with sexually transmitted disease and this could result in severe psychosocial distress shame feelings especially in our culture, 2- upper limbs (site of predilection flexors of wrists and forearm) mainly exposed areas increasing sense of disfigurement, decreased self esteem, more anxiety and depression.

c- Gender (females); mainly because the mean age of our sample was 23.26 ± 9.65 years being at age to have a partner or early marital years; having a dermatological disorder may generate a sense of moving from a beautiful or attractive one to a good or disfigured one

d- Occupational status; those dealing or working with publics or other groups (teachers, nurses...) especially if lichen affecting exposed area or associated with severe itching.

e- Low economic status (may be due to associations with crowdedness which may disturb skin permeability barrier, delaying skin healing in addition to psychosocial and behavioral adverse effects of crowdedness).

f- Psychiatric symptoms (inadequate coping strategies, more psychological stress and more endocrine and immune dysregulations). Our study has several limitations including small size of sample, lack of lichen planus with oral mucosal involvement, lack of correlation of our results with immune studies in skin lesions.

CONCLUSION

Lichen planus is to some extent a psychosomatic or somatopsychic disease or both. Dermatologists should pay attention to the emotional state of their patients knowing that they treat human beings not a skin lesion. Emotional factors are important

as evidenced by higher frequency of psychiatric symptoms, poor quality of life, higher level of anxiety and neuroendocrine and immune dysregulations. Liason consultations between psychiatrist and dermatologist may more helpful in dealing with lichen planus as an autoimmune disorder with no cure but counseling may help in ameliorating symptoms, improving quality of life and enhancing recovery.

REFERENCES

1. Slominski A & Worstman J. Neuroendocrinology of the skin. *Endocr Rev* 2000;21:451-87.
2. Zouboulis CC & Bohm M. Neuroendocrine regulation of sebocytes. A pathogenesis link between stress & acne. *Exp Dermatol* 2004;13(Suppl 4):31-35.
3. Brig Y, Sharma K, Brig S et al. Skin and Psyche ; *Diversitary Symbiosis*, MJAFI 2006;61:163-66.
4. Kimyai-Asadi A & Usman A. The role of psychological stress in skin disease. *J Cutan Med Surg* 2001;140-45.
5. Magin PJ, Pond CD, Smith WT et al. A cross sectional study of psychological comorbidity in patients with acne, psoriasis and atopic dermatitis in specialist dermatology and general practices. *Eur Acad Dermatol Venereol* 2008;23(2):239-40.
6. Koo JYM & Lee CS. General approach to evaluating psychodermatological disorders In Koo JYM , Lee CS .eds., *Psychocutaneous medicine* ,New York NY ; Marceel Dekker Inc, 2004;1-29.
7. Aria N & Kauffman CL. Important drug interactions & reactions in dermatology. *Dermatol Clinics* 2003;21:207-8.
8. Sugerman PB, Savage NW, Walsh Lj et al. The pathogenesis of oral lichen planus. *Crit Rev Oral Biol Med* 2002;13:350-65.
9. Mansur AT, Kilic Z, Atalay F. Psychological evaluation of patients with lichen planus; *Dermatol Psychosom* 2004;5:132-36.
10. Porter SR, Kirby, Olsen I et al. Immunologic aspects of dermal & oral lichen planus. *Oral Surg, Oral Med, Oral Pathol, Oral Radiol Endod* 1997;83:358-66.
11. Manolache I, Seceleanu –Petrescu D, Benea V, et al. Lichen planus patients and stressful events. *J Eur Acad Dermatol Venereol* 2007;22(4):437-41.
12. Koray ML, Dugler O Ak, et al. The evaluation of anxiety and salivary cortisol of oral lichen planus, *Oral Disease* 2003;9(6):298-301.
13. Mignogna MD, Russol L, Fedel S. et al. Clinical behaviour of malignant transforming oral lichen planus. *Eur J Surg Oncol* 2002;28:838-43.
14. Peter ML, Godaert GL, Ballieux RE et al. Moderation of psychological stress responses by personality traits & daily hassles; less flexibility of immune system responses. *Biol Psychol* 2003;65:21-48.
15. Iijima W, Ohtani H, Nakayama. et al. Infiltrating CD8+ T cells in oral lichen planus predominantly express CCR5 & CXCR3 & carry respective chemokines ligands RANTES/CCL5 & IP-10 /CXCL 10 in their cytolytic granule; a potential self recruiting mechanism .*Am J Pathology* 2003;163:261-8.
16. Pipic ML, Kastelan M, Gruber F et al. Perforin Expression in peripheral blood lymphocytes and skin infiltrating cells in lichen planus patients. *Br J Dermatology* 2004;151:433-39.
17. Slominski A & Worstman J. Self regulated endocrine systems in the skin. *Minerva Endocrinol* 2003;28:135-43.
18. World Health Organization ICD-10 Symptoms Checklist for mental disorders version 1.1, Division of Mental Health , Geneva 1994.
19. Ware JE, Snow KK, Kosins KIM et al. SF-36 health survey; manual & interpretation guide. Boston; Health Institute, New England Medical Center 1993.
20. Cohen KR, Demers –Desrosiers LA, Catchlove RFH. A quantitative scoring system for AT9 test as a measure of symbolic function central to alexithymic presentation. *Psychosom* 1938;39:77-88.
21. Beck AT. Beck depression inventory. New York: Harcourt 1996.
22. Taylor AJ. A personality scale of manifest anxiety. *J of Lib Soci. Psychol* 1953;48:285.
23. Aradal E, and Holm AC. Cortisol in saliva reference ranges and relation to cortisol in serum. *Eur J Clin Chem Clin Biochem* 1995;33:927-32.
24. Okumura T, Nakajima Y, Takamatsu T et al. Column switching high performance liquid chromatography system with a laser induced

- fluorimetric detectors for direct, automated assay of salivary cortisol. J. Chromatog B. Biomed. Application 1995;670:11-20.
25. Mohammad J. Psychodermatology: A Guide to understanding Common Psychocutaneous Disorders. Prim Care Companion J Clin Psychiatry 2007;9(3).
26. Shaon M. Disfigurement personal, psychosocial and ethical aspects; J Med. Humanities 2006;110-19.
27. Andrzej S. Neuroendocrine system of the skin, Dermatology 2005;211:199-208.

Address of Correspondence:

ELTawil M E.

Lecturer of Psychiatry, Faculty of Medicine
Cairo University, Cairo, Egypt.

e-mail: Adhamadham3000@yahoo.com

الجوانب النفسية والحيوية لمرضى الحزاز المسطح

تهدف هذه الدراسة إلى محاولة إيجاد مدى التداخل بين الأمراض الجلدية والحالة النفسية، وإرتباط ذلك بتأثيره على جودة الحياة واضطراب الوظائف البيولوجية في الغدد والمناعة. و من هذا النطلق تم دراسة مجموعة تتكون من 70 مريض شخصوا بمرض الحزاز وتم مقارنتهم بمجموعة من 50 مريض ممن يترددون على مركز الأمراض الجلدية. وقد تم استخدام قائمة الأعراض الخاصة بالتصنيف الدولي العاشر للأمراض النفسية ومقياس أس أف - 36 لقياس جودة الحياة ومقياس بيك للاكتئاب وتاييلور للقلق ومقياس عسر المزاج. كما تم أيضا عمل الاختبارات المعملية الخاصة بنسبة الكورتيزون باللعاب و نسبة الخلايا المناعية. أظهرت هذه الدراسة وجود فروق ذات دلالات إحصائية بين المجموعتين في كل المقاييس كما أشارت النتائج أيضا إلى أن وجود بعض العوارض النفسية لمرضى الحزاز يزيد من حدة الإحساس بانحدار جودة الحياة وارتفاع القلق والاكتئاب. كما اكتشف أيضا وجود اضطرابات في المقاييس المعملية. وقد خلصت تلك الدراسة إلى أهمية الاهتمام بالنواحي النفسية والمزاجية عند علاج مرضى الحزاز المسطح خاصة أن هذا المرض ليس له علاج فعال ولكن التدعيم النفسي قد يساعد في تقليل حدة الأعراض وتحسن جودة الحياة عند هؤلاء المرضى.