

Comorbidity of Social Phobia in a Sample of Out-patients with Schizophrenia

El Masry N, Abdel Fattah N, Fouad A

Department of Psychiatry, , Faculty of Medicine, Zagazig University, Sharkia, Egypt.

ABSTRACT

Background: Social phobia in schizophrenia has no enough clinical attention. It has been associated with risk of suicide, poor social functioning and increase risk of relapse.

Objective: to detect and assess social phobia in a sample of out-patients with schizophrenia.

Subjects and methods: Eighty-six patients with schizophrenia and 21 patients with social phobia disorder were diagnosed according to structured clinical interview for DSM-IV. Positive and negative psychotic symptoms in schizophrenia were assessed with scale for assessment of positive symptoms (SAPS) and scale for assessment of negative symptoms (SANS). Social anxiety symptoms were assessed with the Liebowitz Social Anxiety Scale (LSAS). Functioning and disability of the schizophrenia patients were assessed with the medical outcomes study 36-item short form health survey (MOS-36). Results: 22.1% of schizophrenia patients were diagnosed as suffering from comorbid-social phobia, 42.1% of those patients were under treatment by Clozapine. **Results:** There was no significant difference between the schizophrenia patients with comorbid social phobia and without on the SANS and SAPS sum of global scores. Social anxiety scores of schizophrenia patients with comorbid social phobia didn't differ from those with social anxiety disorder as their primary diagnosis. Schizophrenia patients without comorbid social phobia disorder had significantly lower total scores on LSAS and lower social and performance anxiety subscale scores than the other two groups. There were statistically significant differences in functioning and disability on MOS-36 item scale between the two schizophrenic groups. **Conclusion:** Functioning was impaired in patients with comorbid social phobia disorder. Social phobia disorder is not a rare comorbid condition in schizophrenia, and it should be suspected in socially impaired subjects. Its presence implies the need for both psychological and pharmacological therapeutic specificity and often a comprehensive treatment approach.

Key words: Social Phobia, Schizophrenia, Impaired functioning, Egypt.

(Current Psychiatry 2009;16(4):397-402)

INTRODUCTION

Psychiatric comorbidities are common among patients with schizophrenia¹. The presence of anxiety disorders in schizophrenia patients has been associated with a greater risk of suicide², poor social functioning and an increase risk of relapse³.

Schizophrenia and social phobia each cause significant impairment. Social phobia in schizophrenia has no enough clinical attention, although it is important to diagnose and manage social phobia among patients with schizophrenia⁴.

In epidemiological study the rate of social anxiety disorder in schizophrenia range from 13% to 39%^{4, 6-8}.

It is often remains unrecognized and untreated because of its confusion with negative symptoms in schizophrenia patients. Meehl discussed this psychological debate pointing out that anhedonia, a core negative symptom, could contribute to or be a consequence of what he described as "aversive drift in schizophrenia" i.e the tendency to take on a burdensome, threatening, gloomy negative emotional change⁹. He suggested that this aversive drift is intense and pervasive in the interpersonal domain, manifesting itself as ambivalence and interpersonal fear¹⁰.

Social phobia disorder contributes to decreased quality of life in schizophrenia¹¹. This was expected as a comorbid condition causes impairment in quality of life, lower work productivity and greater utilization of health services¹². Those patients also suffer from impairment in occupational role, social relationships and daily life activities¹³.

Social anxiety disorder symptoms are common, quite severe and not correlated with psychotic symptoms severity among schizophrenia outpatients¹⁴.

This study was conducted to determine the prevalence of social phobia in a sample of out-patients with schizophrenia and its effects on functioning of those patients.

SUBJECTS AND METHODS

This study was conducted in Psychiatry Department of Zagazig University Hospitals between June 2008 and February 2009. Patients were selected by using a randomized controlled trial. 86 out-patients diagnosed schizophrenia and 21 out-patients with social phobia disorder according to structured clinical interview for DSM-IV¹⁵ participated in the study.

Informed consents were obtained from the patients. Patients with substance misuse, schizo-affective disorders, neurological disorders and cardiac problems were excluded. All patients were evaluated for clinical symptoms and demographics. The schizophrenia patients were under medical treatment (32: Risperidone; 14: clozapine and 40: typical antipsychotics). 49 schizophrenia patients were males (56.9%), and 37 patients were females (43.1%). Their age range was from 19 to 51 years (mean 28.7±6.3). Schizophrenia patients with comorbid social phobia were 19 patients and without comorbid social phobia were 67 patients. Schizophrenia group with comorbid social phobia included 57.9% males and 42.1 females. Social phobia disorder group included 10 females and 11 males; their age range was from 19 to 45 years.

Positive and negative psychotic schizophrenia symptoms were assessed using the scale for assessment of positive symptoms (SAPS)¹⁶ and the scale for assessment of negative symptoms (SANS)¹⁷.

LSAS¹⁸ was used for schizophrenia patients with comorbid social phobia and patients with social phobia disorder to assess the range of social interaction and performance situations which patients with social anxiety disorder may fear.

Functioning and disability of the schizophrenia patients were evaluated with the medical outcomes study 36-item short-form health survey¹⁹. It includes eight domains (physical function, role-function physical, role-function emotional, social function, mental health, general health, vitality and pain). The domain scores are rated so that higher values indicate better health (range 0-100).

Social anxiety disorder patients were evaluated before the beginning of treatment. Regarding the statistical analysis,

data were collected and analyzed by SPSS statistical software package. ANOVA and post hoc test, T test and chi-square test were used when appropriate. $P < 0.05$ was considered statistically significant.

RESULTS

Table (1) shows that social phobia disorder was diagnosed in 19 (22.1%) schizophrenia patients. No significant difference was found between the two groups of schizophrenia patients with and without social phobia in age, sex and age at start of illness. 42.1% of schizophrenia patients with social phobia disorder were under treatment with Clozapine.

No significant difference was found between the two subgroups on the SANS and SAPS sum of global scores.

Table (2) shows that schizophrenia patients with comorbid social phobia disorder had a significantly higher total score on LSAS (65.7 ± 29.3 vs 48.6 ± 18.8) and a significantly higher scores for total anxiety as well as none significantly higher scores for total avoidance.

Both the performance anxiety and social anxiety scores appeared significantly higher in patients with schizophrenia with comorbid social phobia disorder than in

those without social phobia disorder, whereas no statistically significant differences emerged for performance avoidance and social avoidance scores.

No significant differences were found between schizophrenia patients with social phobia disorder and patients with social phobia disorder as their primary diagnosis in total score on the LSAS (65 ± 29.3 vs 66.6 ± 24.5) or the performance and anxiety scores.

Schizophrenia patients with social phobia disorder and patients with social phobia disorder as their primary diagnosis had significantly higher total scores on the Liebowitz scale than the schizophrenia patients without comorbid social phobia disorder (65.7 ± 29.3 , 66.6 ± 24.5 and 48.6 ± 18.8 respectively).

As seen from table (3) there were significant differences in functioning between schizophrenia patients with and without comorbid social phobia disorder. The affected areas were mentally healthy, role-emotional, social functioning, vitality and general health, whereas no significant differences in physical function and role physical, bodily pain components between the two schizophrenia groups.

Table (1): Demographic and clinical features of the schizophrenia patients.

Variables	Schizophrenia patients group (N=86)			
	Patients with comorbid social phobia (N=19) 22.1%		Patients without comorbid social phobia (N=67) 77.9%	
	N	%	N	%
Sex				
Male	11	57.9	38	56.7
Female	8	42.1	29	43.3
Current treatment				
Clozapine	8	42.1*	6	8.95
Risperidone	6	31.6	26	38.82
Typical antipsychotics	5	26.3	35	52.23
	Mean	S.D	Mean	S.D
Age (years)	28.7	7.3	29.4	6.2
Age at start of illness	19.2	9.3	21.1	6.1
SAPS (sum of global scores)	13.2	3.6	12.9	4.1
SANS (sum of global scores)	15.3	4.8	13.1	4.0

* $P < 0.05$ significant.

Table (2): Liebowitz Social Anxiety Scale scores in schizophrenia patients and in patients with social phobia disorder.

Item	Schizophrenia patients with comorbid social phobia disorder (N=19)		Schizophrenia patients without comorbid without social phobia disorder (N=67)		Social phobia disorder (N=21)	
	Mean	S.D	Mean	S.D	Mean	S.D
Social anxiety	16.1	8.1	10.9	7.1*	16.9	6.8
Performance anxiety	18.1	7.9	11.1	6.8*	19.1	6.2
Social avoidance	16.1	8.1	14.3	7.1	15.8	6.9
Performance avoidance	16.3	9.1	14.6	7.4	17.5	7.1
Total score	65.7	29.3	48.6	18.8*	66.6	24.5

ANOVA for the three groups. *P<0.05 significant when compare with other group.

Higher scores indicate more severe of anxiety symptoms, 55-65 moderate, 65-80 marked, 80-95 severe, >95 very severe

Table (3): Functioning and disability in schizophrenia patients with and without comorbid social phobia disorder on MOS^a.

Item	Schizophrenia patients group (N=86)			
	Schizophrenia patients with comorbid social phobia disorder (N=19)		Schizophrenia patients without comorbid social phobia disorder (N=67)	
	Mean	S.D	Mean	S.D
Physical function	62.2	20	65.3	16.5
Role-physical	58.8	18.2	62.3	24.8
Bodily pain	68.0	24.9	71.6	17.1
General health	59.0	16.1	70.2	19.4*
Vitality	42.3	22.0	53.6	21.6*
Social function	26.5	18.9	57.8	20.3*
Role-emotional	36.7	19.8	54.4	30.6*
Mental health	48.8	19.6	63.2	17.1*

^aMOS: Medical outcomes study 36-item short form health survey: Higher scores indicate better health status (Possible range from 0-100).

*P<0.05 significant.

DISCUSSION

Social phobia disorder constitutes a significant problem for people with psychosis. Schizophrenia and social phobia each cause significant impairment. So this study was conducted to estimate the prevalence of social phobia in a sample of out-patients with schizophrenia and its effects on functioning of those patients.

The study revealed that 22.1% of the schizophrenia outpatients included in this study has a comorbid social phobia disorder. This result is in accordance with Cosoff and Hafner results as they reported that social phobia disorder is the most prevalent anxiety disorder in schizophrenia patients (17%)⁷. Michail and Birchwood examined the rate of social anxiety disorder in first-episode psychosis and reported that 25% of a group of 80 patients fulfilled ICD-10 diagnostic criteria for social anxiety disorder, with a further 11% reported

difficulty in social interaction²⁰. Another study revealed that 11% of those patients were diagnosed as suffering from social phobia⁴. In another study the social anxiety disorder was present in 36.3% of their out-schizophrenic patients²¹.

There is difference between our study result and other results, this may be due to different sample size, different methodology and the collected sample may be from outpatients or inpatients. Low positive and negative symptoms scores on SANS and SAPS were detected in our sample. This is may be due to that our samples are from out-patients, who were already in remission.

Social phobia symptoms become evident when patients start to interact with others and adapt socially and be accepted. There is a similar rate of positive and negative symptoms in both groups of schizophrenia

patients, those with and without comorbid social anxiety disorder. The experience of paranoia and the social withdrawal found in schizophrenia can mimic social anxiety disorder symptoms. Clinical attention to focus on the presence of anxiety symptoms associated with avoidance behavior and on the level of insight, both of which are reduced when negative symptoms are predominant in the clinical picture²². In schizophrenia, withdrawal behavior linked to negative symptoms is phenomenologically sustained by detachment, while social anxiety is related to interpersonal sensitivity.

The social phobia disorder, studied by LSAS¹⁸, was due to stressful situations which need performance for some actions in front of others and not due to paranoia experience. The liebowitz scale for assessment of social anxiety seems adequate and reliable; it was used in similar studies^{4, 21}. This result is accordance with Stern et al. who found that social anxiety disorder symptoms are common severe and not correlated with psychotic symptoms¹⁴. The studied schizophrenia patients with comorbid social phobia disorder, 42.1% of them were under treatment by Clozapine (dose 1000mg in chlorpromazine equivalent).

This is in agreement with studies which found that social anxiety disorder induced by Clozapine treatment and they found that Fluoxetine augmentation improved social anxiety symptoms²³⁻²⁴.

Schizophrenia patients with comorbid social phobia disorder showed low scores of functioning on SF-36 scale items. Affected areas were mental health, emotional-role, social functioning, vitality and general health. All these areas affect quality of life of those patients and leads to impaired functioning, lower productivity and greater utilization of health services. These results are in agreement with results of previous studies^{3-4, 12, 20-21}.

Psychological intervention has been indicated as a possible effective strategy. Group-based cognitive behavior therapy for social phobia in schizophrenia patients improves anxiety symptoms and functioning social¹¹. Particular care is therefore required, when prescribing compounds like Clozapine to schizophrenia patients.

CONCLUSION

Social phobia disorder is not a rare comorbid condition in schizophrenia, and it should be suspected in socially impaired subjects. Its presence implies the need for both psychological and pharmacological therapeutic specificity and often a comprehensive treatment approach.

REFERENCES

1. Buckley PF, Miller BJ, Lehrer DS et al. Psychiatric comorbidities and schizophrenia. *Schizophr Bull* Mar 2009;35(2):383-402.
2. Taiminen T, Huttunen J, Helia H, et al. The schizophrenia suicide risk scale (SSRS): development and initial validation. *Schizophr Res* 2001;47:199-213.
3. Blanchard JJ, Mueser KT, Bellack AS. Anhedonia, positive & negative affect & social functioning in schizophrenia. *Schizophr Bull* 1998;24:413-24.
4. Mazeh D, Bodner E, Weizman R et al. Comorbid social phobia in schizophrenia. *Int J So Psychiatry* May 2009;55:198-202.
5. Bermanzohn PC, Porto L, Arlow PB, et al. Hierachial diagnosis in chronic schizophrenia: a clinical study of co-occurring syndromes. *Schizophrenia Bull* 2000;26:517-25.
6. Cassana GB, Pini S, Saettoni M et al. Multiple anxiety disorder comorbidity in patients with mood spectrum disorder with psychotic features. *Am J Psychiatry* 1999;156:474-76.
7. Cosoff SJ & Hafner RJ. The prevalence of comorbid anxiety disorder in schizophrenia, schizoaffective disorder and Bipolar disorder. *Aust NZJ Psychaitry* 1998;32:67-72.
8. Maria-Michail & Max Birchwood Social anxiety disorder in first-episode psychosis: incidence, phenomenology & relationship with paranoia. *BJP* 2009;195:234-41.

9. Meehl PE. Toward an integrated theory of schizoaffective, schizotypy and schizophrenia. *J Personal Disord* 1990;4:1-99.
10. Meehl PE. Schizoaffective, Schizotypy and schizophrenia. *Am Psychol* 1962;17:827-38.
11. Halperin S, Nathan P, Drummond P et al. A cognitive-behavioral group-based intervention for social anxiety in schizophrenia. *Austr NZJ Psychiatry* 2000;34:809-13.
12. Katzelnick DJ, Koback KA, Deleire T et al. Impact of generalized social anxiety disorder in managed care. *Am J Psychiatry* 2001;158:1999-2007.
13. Wittchen HU, Fuetsch M, Sonntag H, et al. Disability & quality of life in pure & comorbid social phobia-findings from a controlled study. *Eur Psychiatry* 1999;14:118-31.
14. Stern RG, Meiraj H, Ballou S. High social phobia scale scores in schizophrenia don't correlate with psychosis symptom severity scores. Annual Meeting New Research Program and Abstracts. Washington, DC, APA 1999;66.
15. First MB, Spitzer RL, Gibbon M et al. Structured clinical interview for DSM-IV Axis I disorders (SCID), Clinician version. 1995 Washington, DC, American Psychiatric Press.
16. Andreasen N. The scale for assessment of positive symptoms (SAPS). 1986 Iowa city, Iowa.
17. Andreasen N. The scale for assessment of negative symptoms (SANS). 1982 Iowa city, Iowa.
18. Liebowitz MR. Social phobia. *Mod Prol Pharmacopsychiatry* 1987;22:141-73.
19. Mc Hornety CA, Ware JE, Raczek AE. The Mos 36-item short form health survey (SF-36): II-psychometric and clinical tests of validity in measuring physical and clinical tests of validity in measuring physical and mental health constructs medical care 1993;31:93-105.
20. Michail and Birchwood Br. Social anxiety disorder in first-episode psychosis: incidence. *J. Psychiatry* 2009;195:234-41.
21. Pallanti S., Quercioli L., Hollander E. Social Anxiety in Outpatients With Schizophrenia: A Relevant Cause of Disability. *Am J Psychiatry* 2004;161:53-8.
22. Penn DL, Hope DA, Spaulding W et al. Social anxiety in schizophrenia. *Schizophr Res* 1994;11:277-84.
23. Pallanti S, Cuercioli L, Pazzagli A. Social anxiety and premorbid personality in paranoid schizophrenic patients treated with clozapine. *CNS Spectr* 2000;9:29-43.
24. Pallanti S, Cuercioli L, Rossi A et al. The emergence of social phobia during clozapine treatment and its response to fluoxetine augmentation. *J Clin Psychiatry* 1999;60:819-23.

Address of Correspondence:

El-Masry N. Lecturer of Psychiatry
Faculty of Medicine, Zagazig University,
Sharkia, Egypt.
e-mail: nagda.elmasry@yahoo.com

المصاحبة المرضية بين اضطراب الرهاب الاجتماعي

في عينة من مرضى العيادات الخارجية من المصابين بالفصام الذهاني

لا يلقي الرهاب الاجتماعي المشترك لمرض الفصام الذهاني انتباهها الكليين كافيًا، حيث إن له آثارًا تتزامن مع طبيعته و تظهر في شكل محاولات الانتحار وعدم الكفاءة الاجتماعية وسرعة إنتكاسة المرض. وقد أجريت هذه الدراسة لكشف وتقدير اضطراب الرهاب الاجتماعي في عينة من مرضى الفصام الذهاني المترددين على العيادة الخارجية بقسم الطب النفسي بجامعة الزقازيق. شارك في البحث 86 مريضًا بالفصام الذهاني، و 21 مريضًا بالرهاب الاجتماعي تم تشخيصهم حسب مقابلة نفسية طبقًا للتقسيم التشخيصي الإحصائي الأمريكي الرابع للأمراض النفسية. و من ثم تم تقييم الأعراض الذهانية السلبية والإيجابية في مرضى الفصام بواسطة مقياس للأعراض السلبية ومقياس للأعراض الإيجابية. تم تقييم أعراض الرهاب الاجتماعي بواسطة مقياس ليبووترز (LSAS) وتم تقييم الأداء الوظيفي والإعاقات الوظيفية بواسطة (MOS-36). وقد خلصت الدراسة إلى وجود اضطراب الرهاب الاجتماعي في 22.1% من مرضى الفصام (86 مريضًا) ووجد أن 42.1% من مرضى الرهاب الاجتماعي كانوا تحت العلاج بعقار كلوزابين. لا توجد فروق إحصائية بين مقياس الأعراض السلبية والإيجابية بين مرضى الفصام المصاحب له الرهاب الاجتماعي وبين مرضى الفصام بدون الرهاب الاجتماعي. وكانت نتائج مقياس القلق في مرضى الفصام الذهاني المصاحب للرهاب الاجتماعي قياسًا مع مرض الرهاب الاجتماعي كتشخيص أولي متساوية إحصائيًا. وسجل مرضى الفصام بدون الرهاب الاجتماعي أرقامًا أقل من مرضى الفصام المصاحب للرهاب الاجتماعي على مقياس ليبووترز (LSAS). كما كانت الوظائف الاجتماعية والأداء لمرضى الفصام الذهاني المصاحب لمرضى الرهاب الاجتماعي أكثر تأثرًا من مرضى الفصام الذهاني الغير مصاحب لمرضى الرهاب الاجتماعي. لذا نجد أن اضطراب الرهاب الاجتماعي ليس نادرًا بين مرضى الفصام الذهاني ولا بد من تشخيصه. و جدير بالذكر أن هناك ضرورة لإيجاد علاجات سيكولوجية ودوائية لهذه الحالات.